

DG categories (Possibly invented by M. Kelly at UIUC!)
(Ref. Bondal - Kapranov, Keller)

Abelian category: abstraction that captures structure
of $\text{Ab} = \text{category of abelian groups}$
and $R\text{-mod} = \text{cat. of modules / } R,$
 R a ring.

Triangulated category: Abstraction that captures structure
of $K(A) = \text{homotopy category of complexes in } A$
or $D(A) = \text{derived category of } A = K(A)[\text{qiso}^-]$
(A an abelian category)

What about the category $\text{Ch}(A)$ of chain complexes itself?
It is most naturally a Differential Graded (DG) category.

Let k be a commutative ring (often we take k a field)

We say \mathcal{C} is a k -linear category if for any
objects $X, Y \in \text{ob } \mathcal{C}$, the hom set
 $\text{hom}_{\mathcal{C}}(X, Y)$ is equipped w/ struct. of k -module,
and composition

$\text{hom}_{\mathcal{C}}(Y, Z) \times \text{hom}_{\mathcal{C}}(X, Y) \rightarrow \text{hom}_{\mathcal{C}}(X, Z)$
is k -bilinear, allowing us to write it as a map
 $\text{hom}_{\mathcal{C}}(Y, Z) \otimes_k \text{hom}_{\mathcal{C}}(X, Y) \rightarrow \text{hom}_{\mathcal{C}}(X, Z)$

A k -linear category is called additive if it admits
all finite direct sums of objects.

A DG category is a k -linear category \mathcal{C} with the additional structures of

- a \mathbb{Z} -grading $\text{hom}_{\mathcal{C}}(X, Y) = \bigoplus_{i \in \mathbb{Z}} \text{hom}_{\mathcal{C}}^i(X, Y)$ on each hom-module
 - an operator $d_{X, Y}^i: \text{hom}_{\mathcal{C}}^i(X, Y) \rightarrow \text{hom}_{\mathcal{C}}^{i+1}(X, Y)$
- This makes $(\text{hom}_{\mathcal{C}}(X, Y), d_{X, Y})$ into a cochain complex $d^2 = 0$

Such that

- composition $\circ: \text{hom}_{\mathcal{C}}^i(Y, Z) \otimes_k \text{hom}_{\mathcal{C}}^j(X, Y) \rightarrow \text{hom}_{\mathcal{C}}^{i+j}(X, Z)$ is a chain map (commutes w/ differentials)
- $d_{X, X}^0(\text{id}_X) = 0$.

The example is $\text{Ch}(R\text{-mod})$ where R is a k -algebra

Objects = cochain complexes of left R -modules

$$(K^\bullet, d_K) \quad d_K: K^i \rightarrow K^{i+1} \quad d_K^2 = 0$$

R -module hom

$$\text{hom}((K^\bullet, d_K), (L^\bullet, d_L)) = \bigoplus \text{hom}^p((K^\bullet, d_K), (L^\bullet, d_L))$$

$$\text{hom}^p((K^\bullet, d_K), (L^\bullet, d_L)) = \prod_i \text{hom}_R(K^i, L^{i+p})$$

Definition of the space $\text{hom}^p(K, L)$ does not involve d_K or d_L . Elements of $\text{hom}^p((K^\bullet, d_K), (L^\bullet, d_L))$ are not necessarily chain maps.

Differential on $\text{hom}^p((K^\bullet, d_K), (L^\bullet, d_L))$ involves "commuting with the internal differentials"

Formally expect that if $x \in K^i$ and $f \in \text{hom}^j(K, L)$ then $f(x) \in L^{i+j}$

By Leibniz, would have

$$d_L(f(x)) = d_{\text{hom}(K,L)}(f) \cdot x + (-1)^j f(d_K(x))$$

We define $d_{\text{hom}(K,L)}(f) = d_L \circ f - (-1)^{\deg f} f \circ d_K$

These definitions make $\text{Ch}(R\text{-mod})$ into a DG-category

The cohomology or homotopy category of a DG category

Given a DG category \mathcal{C} , we make from there is a subcategory $\mathcal{Z}^0(\mathcal{C})$ with the same objects with morphisms the degree zero cocycles.

$$\text{Hom}_{\mathcal{Z}^0(\mathcal{C})}(X, Y) = \mathcal{Z}^0(\text{hom}_{\mathcal{C}}(X, Y)) = \{f \in \text{hom}_{\mathcal{C}}^0(X, Y) \mid df = 0\}$$

$\mathcal{Z}^0(\text{Ch}(R\text{-mod}))$ is the traditional category of chain complexes and chain maps.

We can also take degree zero cohomology $H^0(\mathcal{C})$

$$\text{Hom}_{H^0(\mathcal{C})}(X, Y) = H^0(\text{hom}_{\mathcal{C}}(X, Y)) = \mathcal{Z}^0(\text{hom}_{\mathcal{C}}(X, Y)) / d \text{hom}_{\mathcal{C}}^{-1}(X, Y)$$

$H^0(\text{Ch}(R\text{-mod}))$ is the traditional homotopy category of chain complexes of R -modules

We can also get a \mathbb{Z} -graded category by considering cohomology of all degrees. $H(\mathcal{C})$

$$\text{Hom}_{H(\mathcal{C})}^i(X, Y) = H^i(\text{hom}_{\mathcal{C}}(X, Y))$$

A DG functor $F: A \rightarrow B$ is a functor that preserves the k -linear structure and direct sums, the grading, and such that

$$F: \text{hom}_A(X, Y) \rightarrow \text{hom}_B(FX, FY)$$

is a chain map.

Such a functor induces $H(F): H(A) \rightarrow H(B)$
 $H^0(F): H^0(A) \rightarrow H^0(B)$

we say F is a quasi equivalence if $H(F)$ and $H^0(F)$ are equivalences of ordinary categories.

That is $F: \text{hom}_A(X, Y) \rightarrow \text{hom}_B(FX, FY)$ is always a quasi-isomorphism of complexes, and $H(F)$ is essentially surjective

If A is a DG category linear over k , then an A -module is a DG-functor $A \rightarrow \text{Ch}(k\text{-mod})$

The collection of all DG functors $\{F: A \rightarrow B\}$ itself forms a DG category $\text{Fun}(A, B)$

Let $F, G: A \rightarrow B$ be two DG functors

Let $\text{hom}_{\text{Fun}(A, B)}^k(F, G)$ be the set of

grading preserving natural transformations $\eta: F \Rightarrow G[k]$, for getting the differential.

So for each $X \in \text{Ob } A$ $\eta_X \in \text{hom}_B^k(FX, GX)$

The differential is just d_B .

Eg $A\text{-mod} = \text{Fun}(A, \text{Ch}(k\text{-mod}))$ is a DG category