

Gradings and Signs on surfaces.

In order to put a \mathbb{Z} -grading on our morphism spaces $\text{hom}(L_0, L_1)$, and also in order to work over a field of characteristic $\neq 2$, we need to make some other choices and restrict the class of curves that we consider.

Let S be a Riemann surface, possibly with boundary.

The first choice we must make is a nowhere vanishing C^∞ quadratic differential

$$\eta_S^2 \in \Gamma(S, (T^*S)^{\otimes 2}) \text{ i.e. } \eta_S^2 : TS \xrightarrow{\mathbb{C}} TS \rightarrow \mathbb{C}$$

e.g. if z is local coordinate in $U \subseteq S$ $\eta_S^2 = f dz^2$, $f \in C^\infty(U, \mathbb{C})$
 f nowhere vanishing

η_S^2 is a complex valued quadratic function on TS

$$\text{so the subset } E_S = \{(p, v) \mid p \in S, v \in T_p S, \eta_S^2(v) \geq 0\}$$

is an \mathbb{R} -subbundle of TS (regarded as a rank 2 \mathbb{R} -vector bundle)
 called the **line field**.

Remarks: The existence of η_S^2 is equivalent to
 $2C_1(S) = 0$ in $H^2(S; \mathbb{Z})$

If S is not closed, this condition is always satisfied.
 If S is closed, say genus $= g$, then this condition is satisfied iff $g=1$. The Fukaya categories of other surfaces cannot be \mathbb{Z} -graded using this method.

Once we have chosen η_s^2 , the definition of the Fukaya category depends on this choice in a nontrivial way. But homotopic choices (connected through space of nowhere-vanishing smooth quadratic differentials) lead to equivalent categories.

The subbundle $\xi_s \subset TS$ is an integrable distribution for dimension reasons.

Proof: let X be a local generating vector field for ξ_s . Any two local sections are of the form fX, gX for some local functions f, g .


$$\text{Then } [fX, gX] = (fX(g) - gX(f)) \cdot X \in \xi_s$$

So $\Gamma(S, \xi_s)$ is closed under Lie bracket, and we may apply the Frobenius theorem.


The (unoriented) foliation that integrates ξ_s is a good way to visualize ξ_s or η_s^2 .

$S = \text{Annulus}$


in \mathbb{C}



$$\eta^2 = dz^2$$



$$\eta^2 = \frac{dz^2}{z^2} = d(\log z)^2$$



$$\eta^2 = \frac{dz^2}{z}$$

(ξ not orientable)

The set of homotopy classes of line fields is a torsor for $H^1(S; \mathbb{Z})$. To see this, consider two nowhere vanishing sections $\eta_1^2, \eta_2^2 \in \Gamma(S, (T^*S)^{\otimes 2})$. The ratio $\eta_2^2/\eta_1^2 \in \Gamma(S, \mathbb{C}^\times)$ is a nowhere vanishing section of the trivial bundle. This may be regarded as a map $S \rightarrow \mathbb{C}^\times$. Since \mathbb{C}^\times is a $K(\mathbb{Z}, 1)$, the homotopy classes of maps are

$$[S, \mathbb{C}^\times] \simeq [S, K(\mathbb{Z}, 1)] \simeq H^1(S; \mathbb{Z}).$$

Now consider a curve $L \subset S$, with tangent bundle $TL \subset TSL_L$. An orientation on L amounts to a choice of (homotopy class of) nowhere vanishing section of TL . But $(TL)^{\otimes 2}$ has a canonical class of nowhere vanishing sections. Thus $(T^*L)^{\otimes 2}$ does as well; let's call it γ_L^2 .

Then γ_L^2 and $\eta_S^2|_L$ are two nowhere vanishing sections of $(T^*SL_L)^{\otimes 2}$, and their ratio may be regarded as a map $\eta_S^2/\gamma_L^2 : L \rightarrow \mathbb{C}^\times$ whose homotopy class

$$[\eta_S^2/\gamma_L^2] \in [L, \mathbb{C}^\times] \simeq H^1(L; \mathbb{Z})$$

Choosing an orientation on L , we get $H^1(L; \mathbb{Z}) \simeq \mathbb{Z}$, so that this class may be interpreted as a number called the **rotation number** of L with respect to η_S^2 and \mathbb{C}^\times . (This is an instance of the more general concept of the Maslov class of a Lagrangian submanifold.)

A curve L such that $[\eta_s^2]_{V_L^2} = 0 \in H^1(L; \mathbb{Z})$ is called **gradable**. Any nonclosed arc is gradable, but a null-homotopic curve is never gradable. Otherwise, gradability depends on the curve and the choice of line field $\xi_s (\eta^2)$.

To say that L is gradable is to say that the subbundles TL and $\xi_s|_L$ in $TS|_L$ are homotopic. We would like to choose a specific homotopy.

One way to think of this is as a bundle over $[0, 1] \times L$.

Let $\pi: [0, 1] \times L \rightarrow L$ be projection.

Def A grading on L is an \mathbb{R} -subbundle $H \subseteq \pi^*(TS|_L)$ such that

$$H|_{\{0\} \times L} = \xi_s|_L \quad \text{and} \quad H|_{\{1\} \times L} = TL .$$

For fixed $p \in L$, $H|_{[0, 1] \times \{p\}}$ is a path of \mathbb{R} -subspaces

of $T_p L$. This may also be regarded as a path

$$H(t, p): [0, 1] \rightarrow \text{RP}(T_p S)$$

$$H(0, p) = \xi_s|_p \quad H(1, p) = T_p L$$

* Let us measure rotation in $\text{RP}(T_p S)$ so that
Closed loops are rotation through $n\pi$ for $n \in \mathbb{Z}$

* Positive sense is determined by orientation of S

Let L_1 and L_2 be two curves with gradings H_1 and H_2
let $p \in L_1 \cap L_2$ be a transverse intersection point.

We can get a path in $T_p S$ from $T_p L_1$ to $T_p L_2$
using $H_2(t, p)$ and $H_1(1-t, p)$.

Let $\Theta \in \mathbb{R} \setminus \pi\mathbb{Z}$ be the total angle through
which this path rotates.

The **absolute index** of P is given by
 $i(p) = \lfloor \Theta/\pi \rfloor + 1 \in \mathbb{Z}$.

Next we consider the problem of signs. This
is a **very tricky** problem and we will use some
surprisingly heavy abstraction to tame it.

Let k be a field. The groupoid of k -lines $\text{Line}(k)$

is the category whose objects are one-dimensional
 k -vector spaces, and whose morphisms are
isomorphisms of such vector spaces.

Every morphism in $\text{Line}(k)$ is invertible, so $\text{Line}(k)$
is a groupoid.

Since any two objects in $\text{Line}(k)$ are isomorphic,
a skeleton for $\text{Line}(k)$ consists of any single
 k -line, say k itself, with all its automorphisms.

This remark shows that $\text{Line}(k)$ is equivalent as a category to the multiplicative group $k^\times = k \setminus \{0\}$ regarded as a category with one object.

We will be particularly interested in $\text{Line}(R)$
We can form a certain quotient $\overline{\text{Line}}(R)$

$$\text{Hom}_{\overline{\text{Line}}(R)}(V, W) = \text{Hom}_{\text{Line}(R)}(V, W)/\sim$$

where if $f, g: V \rightarrow W$ are two isomorphisms of R -lines, we declare $f \sim g$ iff
 $\exists \lambda \in R, \lambda > 0$, such that $f = \lambda g$.

The set $\text{Hom}_{\overline{\text{Line}}(R)}(V, W)$ always has two elements

There is an isomorphism of groups $\text{Hom}_{\overline{\text{Line}}(R)}(R, R) \cong \{\pm 1\}$

The category $\overline{\text{Line}}(R)$ is equivalent to the group $\{\pm 1\} \cong \mathbb{Z}/2\mathbb{Z}$ regarded as a category w/ one object.

Slogan: The signs we are so worried about are "["]just"_"" morphisms in $\overline{\text{Line}}(R)$

k -normalization. We need to relate $\text{Line}(R)$ to our coefficient field k . We construct a functor

$$| \cdot |_k : \text{Line}(R) \rightarrow \text{Line}(k) \quad \text{called } \underline{k\text{-normalization}}$$

Let V be an \mathbb{R} -line. There are exactly two orientations of V , let's call them r_1 and r_2

Define $|V|_k = (k \cdot r_1 \oplus k \cdot r_2) / \langle r_1 + r_2 \rangle$

This is the quotient of the 2-dimensional k -vector space spanned by the orientations, modulo the relation that their sum is zero, so it is a k -line.

Any isomorphism $f: V \rightarrow W$ of \mathbb{R} -lines maps orientations to orientations, and it maps opposite orientations to opposite ones, so it induces a map

$|f|_k: |V|_k \rightarrow |W|_k$ which is an isomorphism of k -lines.

Thus we get a functor $| \cdot |_k: \text{Line}(\mathbb{R}) \rightarrow \text{Line}(k)$

Observe that $| \cdot |_k$ factors through $\overline{\text{Line}}(\mathbb{R})$

$$\begin{array}{ccc} \text{Line}(\mathbb{R}) & \xrightarrow{| \cdot |_k} & \text{Line}(k) \\ \pi \searrow & & \nearrow | \cdot |_k \\ & \overline{\text{Line}}(\mathbb{R}) & \end{array}$$

So $| \cdot |_k$ is really something about orientations and signs.

Remark: In terms of groups, the diagram looks like

$$\begin{array}{ccc} \mathbb{R}^\times & \longrightarrow & k^\times \\ \xrightarrow{x \mapsto \frac{x}{|x|}} & & \nearrow \{\pm 1\} \end{array}$$

Tensor products of lines. $\text{Line}(\mathbb{R})$ is symmetric monoidal under tensor product, and every object V has monoidal inverse $V^* = \text{Hom}_{\text{Vect}}(V, \mathbb{R})$

For each $n \in \mathbb{Z}$ there is an endo functor

$$(-)^{\otimes n}: \text{Line}(\mathbb{R}) \rightarrow \text{Line}(\mathbb{R})$$

Since products of positive numbers are positive, this descends to a functor

$$(-)^{\otimes n}: \overline{\text{Line}}(\mathbb{R}) \rightarrow \overline{\text{Line}}(\mathbb{R})$$

There is also a constant functor $C_{\mathbb{R}}: \overline{\text{Line}}(\mathbb{R}) \rightarrow \overline{\text{Line}}(\mathbb{R})$ that sends every object to \mathbb{R} and every morphism to the identity.

Proposition (Even tensor powers are canonically trivial)

If n is even then $(-)^{\otimes n}$ and $C_{\mathbb{R}}$ are isomorphic as functors $\overline{\text{Line}}(\mathbb{R}) \rightarrow \overline{\text{Line}}(\mathbb{R})$.

Proof let V be an \mathbb{R} -line, and let r_1 and r_2 be the two orientations of V , represented by vectors v_1 and $v_2 = -v_1$ respectively.

Because n is even, $v_1^{\otimes n} = v_2^{\otimes n}$ in $V^{\otimes n}$

Let $\eta_V \in \text{Hom}_{\overline{\text{Line}}(\mathbb{R})}(V^{\otimes n}, \mathbb{R})$ be the equivalence class of isomorphisms taking $v_1^{\otimes n} = v_2^{\otimes n}$ to $1 \in \mathbb{R}$

Then η is a natural isomorphism from $(-)^{\otimes n}$ to C_R \square

Proposition If n is odd then $(-)^{\otimes n}$ is isomorphic to the identity functor.

Now let L_1 and L_2 be two graded curves that intersect transversely at p . We have defined an absolute index $i(p) \in \mathbb{Z}$.

If we regard p as a morphism from L_1 to L_2 we can define the **orientation line at p**

$$o(p) = (T_p L_2)^{\otimes i(p)} \text{ as object of } \overline{\text{Line}}(R)$$

That is, we take the tangent space to the target object (L_2) raised to the index.

If $i(p)$ is even then $o(p) \cong R$ canonically.
 If $i(p)$ is odd then $o(p) \cong T_p L_2$ canonically.


The proper definition of $\text{hom}(L_1, L_2)$ is this:
 For transversely intersecting graded curves L_1 and L_2
 There is a graded k -vector space

$$\text{hom}^i(L_1, L_2) = \bigoplus_{n \in \mathbb{Z}} \text{hom}^n(L_1, L_2)$$

$$\text{hom}^n(L_1, L_2) = \bigoplus_{\substack{p \in L_1 \cap L_2 \\ i(p)=n}} |o(p)|_k = \bigoplus_{\substack{p \in L_1 \cap L_2 \\ i(p)=n}} |(T_p L_2)^{\otimes n}|_k$$

This affects the definition of μ^d as follows.

μ^d counts disks, and the boundary orientation of the disk induces orientation on the curves (at least locally near an intersection point)



"apply boundary orientation of disk to target object."

$$\begin{aligned} \text{hom}(L_0, L_1) &\geq |\alpha(p_1)|_k - |(T_{p_1} L_1)^{i(p_1)}|_k \\ \text{hom}(L_1, L_2) &\geq |\alpha(p_2)|_k \\ \text{hom}(L_2, L_3) &\geq |\alpha(p_3)|_k \\ \text{hom}(L_3, L_4) &\geq |\alpha(p_4)|_k \\ \text{hom}(L_0, L_4) &\geq |\alpha(p_0)|_k - |(T_{p_0} L_4)^{i(p_0)}|_k \end{aligned}$$

A disk with inputs p_1, \dots, p_d and output p_0 thus gives rise to a unique morphism in $\overline{\text{Line}}(\mathbb{R})$

$$\alpha(u): \alpha(p_d) \otimes \alpha(p_{d-1}) \otimes \cdots \otimes \alpha(p_1) \rightarrow \alpha(p_0) \quad (\text{match boxes from picture above})$$

The k -normalization is the contribution to μ^d

$$|\alpha(u)|_k : |\alpha(p_d) \otimes \cdots \otimes \alpha(p_1)|_k \longrightarrow |\alpha(p_0)|_k$$

$$|\alpha(p_d)|_k \otimes \cdots \otimes |\alpha(p_1)|_k \quad \cap \quad \text{hom}(L_0, L_d)$$

$$\text{hom}(L_{d-1}, L_d) \otimes \cdots \otimes \text{hom}(L_0, L_1) \longrightarrow$$

It is also a fact that the polygon is rigid iff $i(p_0) = i(p_1) + \cdots + i(p_d) + 2 - d$